



SPECTRUM OF VENTRAL HERNIA IN PATIENTS PRESENTING AT INDIRA GANDHI MEDICAL COLLEGE SHIMLA

Dr. Vipul Parmar¹, Dr. RS Jhobta², Dr. Geeti Parmar³, Dr Vishal Thakur and Dr. Ramesh Kaundal^{4*}

¹Medical officer, MS General Surgery, CH Theog

²Professor Department of General Surgery, IGMC Shimla

³Junior Resident Department of Pharmacology, IGMC Shimla

⁴Assistant Professor Department of General Surgery, SLBSGMC Ner Chauk

⁵Medical Officer, MS General Surgery, CH Theog

ARTICLE INFO

Article History:

Received 06th May, 2020

Received in revised form 14th June, 2020

Accepted 23rd July, 2020

Published online 28th August, 2020

Key words:

Spectrum, Ventral hernia, Incisional hernia, LVHR, Anatomical repair

ABSTRACT

Background: ventral hernia is one of the common complications encountered in surgery OPD day by day and is an important cause of morbidity. It can be repaired by following anatomical, mesh or laparoscopic methods. The incidence of these hernias is high even with recent advances in surgery, anaesthesiology, antibiotics, and suture materials used. We wanted to study the epidemiology, aetiology, mode of presentations, modalities of treatment and its outcome of ventral hernia repair.

Method: with the aim of studying spectrum of the ventral hernia in Indira Gandhi medical college, study was conducted on 207 patients who presented in IGMC Shimla with in two year period. History, clinical examination, radiological as well as laboratory investigations, comorbid conditions, operative techniques, post operative period was recorded on a preformed performa and analysed.

Results: 207 patients were studied age ranging from 10-88 with mean age was found to be 51.33 years with female predominance with ratio 1.5:1. Most of the patients who present in IGMC are from shimla district and Epigastric hernia (41.55%) was commonest variety followed by incisional hernia (30.43%), umbilical, paraumbilical, recurrent incisional and spigelian hernia. 104 patients were symptomatic and prostatism (29.80%) was the most common comorbidity followed by hypertension, respiratory and diabetes. Majority of the patients underwent anatomical repair (37.68%), open mesh hernioplasty 31.67% and laparoscopic ventral hernia was done in 19.8%. Patients followed for one month after surgery for post op complications and only 10 patients (4.8%) of operated patients had wound infection.

Conclusion: Ventral hernias are common surgical burden in Himalayan population. Better treatment strategies need to be devised to manage these. Presence of ventral hernia is an indication for surgery even in presence of co-morbid conditions like ascites, COPD, BPH as these patients are more prone for complications and of course these conditions need proper addressal.

Copyright©2020. Dr. Vipul Parmar et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Ventral Hernia is defined as a protrusion of an abdominal viscus or part of a viscus through the anterior abdominal wall. The term ventral hernia refers to the hernias of the anterior abdominal wall except groin hernias. It represents a heterogeneous subtype of hernia disease that includes incisional hernias, paraumbilical hernias, umbilical hernias, epigastric hernias and spigelian hernias respectively¹Epigastric hernia² is the protrusion or herniation of extraperitoneal fat

through a defect in the linea alba anywhere between the xiphoid process and the umbilicus, usually midway between these structures. This type is typically infrequent in children and is usually found in adults with a male to female ratio of around 3:1.

Incisional hernia² is one where the peritoneal sac herniates through an acquired scar in the abdominal wall usually caused by a previous surgical operation or an accidental trauma.. These are infrequent under the age of 40 years and their incidence increases with age. Ventral incisional hernia comes out to be a common complication of abdominal surgery, with the overall incidence ranging from 2% to 20%. It is very common in females.

*Corresponding author: Dr. Ramesh Kaundal

Assistant Professor Department of General Surgery, SLBSGMC Ner Chauk

Spigelian hernia² comprises about 0.12% of all the abdominal wall hernias. This is a type of inter-parietal hernia occurring at the level of arcuate line through spigelian point. Spigelian hernia can occur above (10%) or below (90%) the umbilicus. Below the umbilicus it occurs at the junction of linea semilunaris and linea semicircularis. Spigelian hernias are most frequently found in adults from ages 40 to 70, typically in the patients around 50 years of age and the male to female ratio is 1:1.8. The incidence of Spigelian hernias in children is very low.

Lumbar hernia² is herniation either through superior (Grynfelt's) or inferior lumbar triangle (Petit's triangle) Approximately 20 per cent of lumbar hernias are congenital while the rest of these are acquired (80%), either primarily or secondarily (25%). Ventral hernias are among the most common clinical problems seen by a surgeon.

Epigastric hernia² is the protrusion or herniation of extraperitoneal fat through a defect in the linea alba anywhere between the xiphoid process and the umbilicus, usually midway between these structures. This type is typically infrequent in children and is usually found in adults with a male to female ratio of around 3:1. Only a few studies are available on the incidence of epigastric hernias, where they account for approximately 1–5% of abdominal wall hernias.

A thorough understanding of the pathophysiology and complete spectrum of ventral hernias can improve management of such patients. Multiple important factors are related to the occurrence and successful repair of ventral hernias which include patient-specific characteristics, presence or absence of comorbidities, technique of repair and the type of material used among others. A very few studies relate to study the spectrum of ventral hernias in the state of Himachal Pradesh in India.

Thus present study aims to find the incidence and prevalence of ventral hernias in the population of Himachal Pradesh which represents a distinct geographic, environmental and diet consuming entity than the rest of the Indian subcontinent. This study also aims to find the effect of various factors including type of repair, choice of material used to repair, presence or absence of associated comorbidities, age and gender on the management and outcomes of ventral hernia repair.

METHODS

The present study is an observational type of study conducted prospectively, in the Department of Surgery at I.G.M.C Shimla, on 207 patients who presented with symptoms of ventral hernia in the Department of Surgery from 1st July 2016 to 30th June 2018.

Inclusion criteria

The following patients were included in the study

- USG proven ventral hernias, admitted in the department of Surgery
- Verbal and written consent was taken from all the patients of ventral hernia and all those who agreed for procedure were included in the study.

Exclusion criteria

The following patients were excluded

- Groin hernia and lumbar hernia diagnosed patients.
- Patients who were reviewed on outpatient basis were excluded

All the patients who met the inclusion criteria evaluated thoroughly and their history, clinical examination and various investigations are recorded in a preformed performa and entered on masterchart in MS excel. Patients were diagnosed and treated accordingly. The patients were evaluated on the following parameters:

- History and Physical examination
- Age
- Gender
- Geographical distribution (district wise)
- Fitness for surgery
- Smoker or non smoker
- Co-morbidities
- Diagnosis
- USG Finding
- Treatment given
- Use of mesh
- Content of hernial sac
- Length of hospital stay
- 1 month follow-up for wound sepsis.

Various treatment options included anatomical repair, open mesh hernioplasty (onlay, sublay and underlay) and laparoscopic ventral hernia repair

All the parameters will be noted as per Performa attached and the data so collected was analyzed statistically using softwares Epi info and latest version of SPSS. P value < 0.05 was considered to be statistically significant.

RESULTS

Over a period of one year, 308 cases of abdominal wall hernias were studied at our institute out of which 16 patients were not operated due to unfit for surgery. Age varies from 10 months to 88 years with an increasing incidence with age (**Table 1**). The mean age of patients presented in this study was found to be 51.33 years. The male to female ratio is 1:1.5 as shown in **Table 2**. Ventral hernias showed a female dominance (**Table 2**). Epigastric hernia was commonest variety followed by incisional hernia, umbilical, paraumbilical, recurrent incisional and spigelian hernia distribution is shown in **table no. 3**. Patients presenting at IGMC Shimla were from all region of Himachal Pradesh and most of the patients are from district Shimla, 97 out of 207 (46.86%) and lowest from Una i.e. only 1 patient (0.48%). Area wise distribution is depicted in table no.4.

104 patients were symptomatic like pain, tenderness, and features of intestinal obstruction. Prostatism was commonest associated illness followed by hypertension and respiratory diseases (**Table 5**). Wound infection was commonest complication in post operative period followed by pyrexia

despite routine use of antibiotics in 10 cases out of 207 patients (4.8%).

After assessing and diagnosing the patients various surgical techniques were performed according to their diagnosis and size of defect. Various approaches are mentioned in table no.6.

Table no 1 Age wise distribution

| Age group (yrs) | No. of patients | % of total patients |
|-----------------|-----------------|---------------------|
| <15 years | 3 | 7.18 |
| 15 to 30 years | 13 | 9.68 |
| 31 to 45 years | 59 | 10.93 |
| 46 to 60 years | 71 | 17.50 |
| >60 years | 61 | 19.06 |

Table no 2 Sex distribution

| Gender | No. of patients | % of total patients |
|--------|-----------------|---------------------|
| Male | 80 | 38.65% |
| Female | 127 | 61.35% |
| Total | 207 | 100.00% |

Table no 3 Typewise distribution of abdominal wall hernia

| Type of hernia | No. Of patients | % of total patients |
|-----------------------------|-----------------|---------------------|
| Epigastric hernia | 86 | 41.55% |
| Para-umbilical hernia | 21 | 10.14% |
| Incisional hernia | 63 | 30.43% |
| Umbilical hernia | 25 | 12.08% |
| Recurrent incisional hernia | 10 | 4.83% |
| Spigelian hernia | 2 | 0.97% |

Table no 4 Area wise distribution

| District | No. of patients | % of total patients |
|----------|-----------------|---------------------|
| Shimla | 97 | 46.86% |
| Mandi | 44 | 21.26% |
| Bilaspur | 14 | 6.76% |
| Kangra | 3 | 1.45% |
| Sirmour | 14 | 6.76% |
| Kullu | 12 | 5.80% |
| Hamirpur | 3 | 1.45% |
| Ludhiana | 1 | 0.48% |
| Solan | 16 | 7.73% |
| Kinnaur | 2 | 0.97% |
| Una | 1 | 0.48% |
| Total | 207 | 100.00% |

Table 5 Comorbid illnesses

| Diseases | No. of patients | % of total patients |
|--------------------|-----------------|---------------------|
| Diabetese Mellitis | 22 | 10.62 |
| Hypertension | 50 | 24.03 |
| Respiratory | 30 | 14.42 |
| Prostatism | 62 | 29.80 |

Table no.6 Surgical Approaches

| Surgery | No. Of patients | % of total patients |
|------------------------------------|-----------------|---------------------|
| Not operated as unfit for surgery | 17 | 8.21% |
| Anatomical repair | 78 | 37.68% |
| Onlay mesh hernioplasty | 66 | 31.88% |
| Laparoscopic ventral hernia repair | 41 | 19.8% |
| Sublay mesh hernioplasty | 3 | 1.45% |
| Underlay mesh hernioplasty | 2 | 0.97% |
| Total | 207 | 100.00% |

DISCUSSION

Being a commonly performed general surgical operation, abdominal wall hernia comprises a significant proportion of total surgical work load in most of the centres³. In present study female patients have the higher incidence of ventral hernia(i.e.61.35%) as compared to males (38.65%) which is in accordance with the study of McAdam Eccles *et al*⁴ and Fanklin *et al* study⁵. Our study shows that maximum incidence of the ventral hernia is in the age group of 31 to 45 i.e. 34.30% and mean age of the patient was found 51.33 year which is similar to the various studies conducted worldwidw^{5,6}. Most common ventral hernia in our study is epigastric hernia followed by incisional hernia and varies in all ages. However this result differ siltly from the study coduncted by sangwan *et al*⁷ where they found paraumbilical hernia as with highest frequency. Epigastric hernia are not uncommon and have been documented as an acquired disease in more than 90% cases⁸. In our study epigastric hernia account for 41.55%. Multiparity, obesity, malnutrition and raised intra abdominal pressure have been assigned as predisposing factors. Incisional hernia was noticed in approximately 30.43% of the cases which is significantly highr than USA and UK where it has been reported in 6% - 10% of cases but in accordance with African literature^{9,10}. At this institute patient presents from all over the himachal Pradesh but maximum no. of the the patients were from shimla district i.e. 46.86% due to nearby area and less no. Of patients are from border area of the himachal because proximity to the other care centres outside the state. Concomittent medical disease has been reported as having role in its etiology as well as in increasing morbidity and mortility in postoperative period. Symptoms of prostatism were present in approximately 29.9% cases while hypertension and diabetes were present in approximately 24% and 10% cases respectively. In the past decades the complication rates of abdominal wall hernias have drastically improved with the use of prosthetic mesh. We have used mesh unanimously in all cases except pediatric, strangulated and small ventral hernias. No recurrence has been reported till date. Complication rate in the form of post operative pain, seroma formation has been noticed. Wound sepsis of 4.8% has been noticed in our study which is in the range of 4% - 12% mentioned in literature¹¹. Complications of the ventral hernia like irreducibility, obstruction and strangulation has been reduced in recent era as increased awareness of population.

Various surgical approaches has been used according to patients diagnosis and defect size. 17 patients were unfit for surgery and rest 190 patients underwent for surgery. Laparoscopic ventral hernia was performed in 19.8% of patients and open hernia repair was done in 80.1% patients. Open mesh herniopalsty was done in 34.3% patients and 37% patients underwent anatomical repair due to their small size defect. Laparoscopic incisional hernia repair is a widely used and accepted operative technique, assuming general advances of laparoscopy are also valid for this group. In our study laparoscopic ventral hernia repair has given promising result as recent studies have shown that in the short term laparoscopic repair is superior to open repair in terms of less blood loss, fewer perioperative complications, and shorter hospital stay.^{12,13}

CONCLUSION

Ventral hernias are common surgical burden in Himalayan population. Better treatment strategies need to be devised to manage these. Presence of ventral hernia is an indication for surgery even in presence of co-morbid conditions like ascites, COPD, BPH as these patients are more prone for complications and of course these conditions need proper addressal. Options for the surgical management of ventral hernias continue to evolve and laparoscopic ventral hernia repair is the treatment of choice in present era.

References

1. Townsend R, Beauchamp B, Mattox M. Clinical surgery of hernia. Sabiston Textbook of Surgery, 19th Edition, Volume II, Elsevier. 2016:1128.
2. Kingsnorth AN, LeBlanc KA, editors. Management of abdominal hernias. Springer Science & Business Media; 2013.
3. Jennings WK, Anson BJ, Wright RR. A new method of repair for indirect Inguinal Hernia considered in reference to parietal anatomy. *Surgery Gynecology Obstetrics*. 1942; 74: 697.
4. Eccles WM. Hernia; Its Etiology, Symptoms, and Treatment. Baillière, Tindall and Cox; 1900.
5. Franklin M, Gonzalez J, Glass J, Manjarrez A. Laparoscopic ventral and incisional hernia repair: an 11-year experience. *Hernia*. 2004;8(1):23-7.
6. Danzig M, Stey A, Yin S, Qiu S, Divino C. Patient profiles and outcomes following repair of irreducible and reducible Ventral Wall Hernias. *Hernia : the journal of hernias and abdominal wall surgery*. 2015.
7. M. Sangwan *et al.* / *Open Journal of Epidemiology* 3 (2013) 135-138
8. Olmi S, Scaini A, Cesana GC, Erba L, Croce E. Laparoscopic versus open incisional hernia repair: an open randomized controlled study. *Surg Endosc*. 2007;21(4):555-55917364151
9. Sultan, B., Qureshi, Z. and Malik, M.A. (2009) Frequency of external hernias in Ayub teaching hospital Abbottabad. *Journal of Ayub Medical College Abbottabad*, 21, 57-58. [9]
10. Garba, E.S. (2000) The pattern of adult external abdominal hernias in Zaria. *Nigerian Journal of Surgical Research*, 2, 12-15.
11. Mbah, N. (2007) Morbidity and mortality associated with inguinal hernia in northwestern Nigeria. *West African Journal of Medicine*, 26, 288-292.
12. Ammar, S.A. and Ismail, T. (2008) Abdominal wall hernias in upper Egypt: A different spectrum. *East and Central African Journal of Surgery*, 13, 109-113.
13. Osifo, O.D. and Amusan, T. (2010) Outcome of giant inguinocrural hernia repair with local lidocaine anaesthesia. *Saudi Medical Journal*, 31, 53-58.

How to cite this article:

Dr. Vipul Parmar *et al* (2020) ' Spectrum of Ventral Hernia in Patients Presenting at Indira Gandhi Medical College Shimla', *International Journal of Current Advanced Research*, 09(08), pp. 22942-22945.
DOI: <http://dx.doi.org/10.24327/ijcar.2020.22945.4535>
