



Research Article

STUDY TO ASSES THE MAJOR PREOPRATIVE PREDICTORS FOR DIFFICULT LAPAROSCOPIC CHOLECYSTECTOMY

Dr Ramesh kaundal¹, Dr Sanjeev Gupta², Dr Seema^{3*}, Dr Vishal Thakur and ⁴Dr Vipul Parmar⁵

¹M.B.B.S,M.S. General Surgery, Medical Officer, C.H.Theog

²M.B.B.S,M.S. General Surgery Associate Professor IGMC Shimla

³M.B.B.S,M.S.Ophthalmology, medical officer DDUZH Shimla

⁴M.B.B.S,M.S. General Surgery, Assistant Professor, SLBSCM Ner Chowk

⁵M.B.B.S,M.S. General Surgery, Medical Officer, C.H.Theog

ARTICLE INFO

Article History:

Received 6th May, 2020

Received in revised form 15th June, 2020

Accepted 12th July, 2020

Published online 28th August, 2020

Key words:

Clinical, Sonological, Difficult laparoscopic cholecystectomy, Predictors

ABSTRACT

Background: Laparoscopic cholecystectomy (LC) has become the gold standard treatment for gallstone disease. Though mostly safe occasionally it can be difficult due to various problems faced during surgical procedure. Anticipation of likely difficulty can help in avoiding complications and legal aspect.

Methods: With the aim of identifying, comparing various predictors of difficulty and their correlation with likely difficulty this prospective study on 100 patients undergoing laparoscopic cholecystectomy for symptomatic cholelithiasis was undertaken. Various clinical, sonological predictors were recorded and assessed for intraoperative difficulties and frequency of difficult laparoscopic cholecystectomy was recorded.

Results: 100 patients were studied with age ranging from 10-80 years, maximum incidence of difficult LC (36.14%) being in the age group 41-50 years. The time taken for LC increased significantly with increasing age. Gender affects the difficulty i.e. 48.15 % difficulty in male as compared to 24.66% among female patients. Body mass index (BMI) > 25 did not show any significance with difficult cholecystectomy. Previous history of multiple attacks of acute cholecystitis, h/o ERCP with CBD stenting, Upper abdominal surgery, clinically tender palpable GB, USG evidence of contracted/distended GB, impacted stone at GB neck, GB wall thickness >4 mm, multiple stones in GB had increased rate of difficult cholecystectomy and the conversion rate.

Conclusions: Clinical and sonological predictors are most reliable factors. Use of good clinical judgment regarding possibility of and likely difficulty along with understanding of available resources is important in making decision in each case.

Copyright©2020. Dr Ramesh kaundal et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Laparoscopic cholecystectomy is the treatment of choice for gall bladder stone disease. Difficult laparoscopic cholecystectomy (LC) is associated with serious complications and a high conversion rate¹. The prevalence of cholelithiasis is 10-15% in India, and approximately 1-2% of asymptomatic patients will develop symptoms and required cholecystectomy every year.² Laparoscopic cholecystectomy has become the procedure of choice for management of symptomatic gallstone disease for its minimal invasive, less pain and early recovery^{3,4}. A number of conditions can be associated with increased difficulty of cholecystectomy. These include acute cholecystitis, severe chronic cholecystitis, over distended due to impacted stone at Hartman pouch or contracted gallbladder,

intraoperative adhesions due to previous upper abdominal surgery, obesity, male gender and post ERCP and other other gall bladder related conditions. With these predisposing factors at times, it takes longer time even with bile/ stone spillage and occasionally it requires conversion to open cholecystectomy. However, of all Laparoscopic cholecystectomies, 1-13% requires conversion to an open for various reasons.⁵

Preoperative recognition and awareness as well as modifications of technical strategies are important to a successful outcome and avoiding major complications in these patients⁶. Assessing patient clinically and sonologically preoperatively for difficult laparoscopic cholecystectomy, and finding out the major factor would be useful for informing patients and a more experienced surgical team could be assembled when risk for conversion appears significant.

*Corresponding author: Dr Seema

M.B.B.S,M.S.Ophthalmology, medical officer DDUZH Shimla

METHOD

All cases of symptomatic gallstones undergoing elective LC at IGMC hospital were studied over one year in a prospective manner.

Inclusion criteria: All the patients undergoing elective laparoscopic cholecystectomy due to following gall stone diseases:

1. Symptomatic cholelithiasis
2. Empyema gall bladder
3. Mucocoele gall bladder
4. Gall bladder polyp
5. Biliary pancreatitis

Exclusion criteria

1. Patients scheduled for Emergency LC for calculous acute cholecystitis or indicated for a primary open procedure.
2. Patients with common bile duct calculus
3. Patient with suspicion of ca gall bladder
4. Features of obstructive jaundice
5. Cholelithiasis during pregnancy
6. Absolute contraindications to LC like cardiovascular, pulmonary disease, coagulopathies and end stage liver disease.
7. Laparoscopic cholecystectomy performed with other laparoscopic intervention in same setting.

The selected patients were counseled about the procedure and written informed consent was taken regarding participation in the study as well as for the surgical procedure. The patients were also informed of the likelihood of conversion to OC. The preoperative work up of the study population involved detailed history taking, clinical examination, mentioning predictors for difficult cholecystectomy on history sheet laboratory investigations and ultrasound of abdomen was done. Proven predictors for difficult LC taken were following: age, body mass index (BMI), whether gallbladder was palpable per abdominally and if tenderness was present in the right hypochondrium. Transabdominal ultrasound was done routinely in the preoperative work up to assess number and size of gallstones in gallbladder, if the gallstones were impacted in gallbladder lumen, if the gallbladder was contracted, the thickness of gallbladder wall >4 mm, wall echo sign and to asses for pericholecystic fluid.

The standardized 4-port technique (2 × 10 mm, 2 × 5 mm) of laparoscopic cholecystectomy is used; the patient and operating team being positioned in the American style. Monopolar electrocautery is used as the haemostatic modality. Pneumoperitoneum was created with carbon dioxide by using veress needle. Considering previous studies from the published literatures

Laparoscopic cholecystectomy is considered as difficult if any of the following 4 criteria is fulfilled during the surgery

1. Time taken from skin incision to skin closure more than 90 minutes.
2. Time taken for Callot's triangle dissection more than 20 minutes

3. Time taken for Gall bladder dissection from gall bladder fossa more than 20 minute.
4. Conversion to open cholecystectomy due to any reason.

Data collected in proforma is transferred into MS Excel sheet for further processing and analysis. Data is further analyzed using statistical software Epi info version 4 and SPSS version 20. Qualitative variables are expressed in term of frequencies, proportion and 95% Confidence Interval while quantitative variables are expressed as mean and standard deviation. In order to compare results between two study groups appropriate parametric or non parametric test of statistical significance are used. Probability value (p-value) less than 0.05 are considered statistically significant.

RESULTS

Hundred (100) patients of Ultrasonographically proved symptomatic cholelithiasis; admitted in a single unit of the surgical wards of department of general Surgery IGMC Shimla; were included in the present study. These patients were admitted from July 2016 through June 2017. Laparoscopic cholecystectomy was performed in all these selected 100 patients among whom 31 patients underwent difficult laparoscopic cholecystectomy; while 69 patients had easy laparoscopic cholecystectomy (figure 1).




Figure 1 Difficulty Level

The study included a total patients with age ranging from 10 years to 78 years with mean age 46.69 +/- SD 13.79 years. The maximum incidence (30%) was seen in the age group of 51 to 60 years (Figure 2) and maximum difficulty was seen in age group 41-50 (48.3%). There were a higher proportion of females (71%).




Figure 2 Age Distribution

In our study 10 patients were operated during acute attack of cholecystitis and 3 out of them faced difficulty in LC and did not showed any significant correlation with difficult cholecystectomy. Multiple attacks of acute cholecystitis was present in 63 patient and 25 (39.7%) showed difficulty. H/O previous upper abdominal surgery was present in 10 patients and 7 (70%) had difficult cholecystectomy and was also reason for conversion. 12 patients had the history of ERCP with CBD stenting and 10 patients had difficult LC. On clinical examination 27 patients had BMI >25 and 7 patients encountered difficulty and 20 LC was easy. Patients with the tenderness in right hypochondrium was present in 7 patients and 5 patents showed significant difficulty while performing LC and was also reason for conversion. On clinical examination palpable Gall bladder was present in 14 patients and was statistically significant for difficult cholecystectomy. On sonological examination contracted gall bladder was present in 37 patients and out of these 64 %,24 patients had difficult cholecystectomy, distended gall bladder was present in 16 patients and was also a strong predictor for difficult cholecystectomy and 11 patient had difficult cholecystectomy. In acute attack of cholecystitis gallbladder was found inflamed in 6 cases on USG and 5 patients associated with difficult cholecystectomy. Wall echo sign was seen in 14 patients on USG and only 4 patients had difficult cholecystectomy and 10 patients had easy cholecystectomy. Gallbladder studded with multiple stone was present in 79 patients and out of these 30 LC was difficult to perform whereas big impacted stone ant neck of gall bladder was present in 24 patients, out of these patients 17 LC found to be difficult. Thick wall GB i.e. GB wall thickness >4 mm was found on 15 cases and out of these cases, 13 LC was difficult. pericholecystic fluid was present in 6 patients and, 4 LC was easy and only 2 were found to be difficult. Predictive association between risk factors and intraoperative outcome with statistical analysis is depicted in table 2.

Table 2 Association between risk factors and intraoperative outcome

Predictors		Essy	Difficult	Total p value
		cholecystectomy	cholecystectomy	
Clinical	Age			
	<30	10 (100%)	0 (0%)	10
	31-40yrs	19 (94.95%)	1 (5.05%)	20
	41-50 yrs	15 (51.7%)	14 (48.3%)	29
	51-60yrs	18 (60.0%)	12 (40.0%)	30
Gender	>60 yrs	7 (63.6%)	4 (36.4%)	11
	Female	55 (75.44%)	18 (24.66%)	73
	Male	14 (41.85%)	13 (48.15%)	27

Acute cholecystitis	Yes	7 (70.0%)	3 (30%)	10	1.00	
	No	62 (68.8%)	28 (31.2%)	90		
Multiple attacks	Yes	38 (60.3%)	25 (39.7%)	63	0.015	
	No	31 (83.8%)	6 (16.2%)	37		
Previous abdominal sugery	Yes	3 (30%)	7 (70.0%)	10	0.019	
	No	66 (73.3%)	24 (26.7%)	90		
ERCP with Stenting	Yes	2 (16.67%)	10 (83.33%)	12	0.01	
	No	67 (83.34%)	21 (16.66%)	88		
BMI	>25	20 (74.1%)	7 (25.9%)	27	0.629	
	<25	49 (67.1%)	24 (32.9%)	73		
Tenderness in RHC	Yes	2 (28.6%)	5 (71.4%)	7	0.02	
	No	67 (72.1%)	26 (27.9%)	93		
Palpable Gall Bladder	Yes	3 (21.4%)	11 (78.6%)	14	0.01	
	No	66 (76.7%)	20 (23.3%)	86		
Contracted GB	Yes	13 (35.1%)	24 (64.9%)	37	0.01	
	No	56 (64.9%)	7 (11.1%)	63		
Distended GB	Yes	5 (31.3%)	11 (68.7%)	16	0.01	
	No	64 (76.2%)	20 (23.8%)	84		
Inflamed GB	Yes	1 (16.7%)	5 (83.3%)	6	0.01	
	No	68 (70.2%)	26 (29.8%)	94		
Wall Echo Sign	Yes	10 (71.4%)	4 (28.6%)	14	0.005	
	No	61 (70.9%)	25 (29.1)	86		
Sonological predictors	Multiple stones	Yes	49 (62.0%)	30 (38.0%)	79	0.003
	No	20 (95.2%)	1 (4.8%)	21		
Impacted Sone	Yes	7 (29.2%)	17 (70.8%)	24	0.001	
	No	62 (81.6%)	14 (18.4%)	76		
GB wall thickness >4 mm	Yes	2 (13.3%)	13 (86.7%)	15	0.001	
	No	67 (78.2%)	18 (21.2%)	85		
Pericholecystic fluid	Yes	4 (66.7%)	2 (33.3%)	6	1.000	
	No	65 (69.1%)	29 (30.9%)	94		

DISCUSSION

In this study, 100 laparoscopic cholecystectomies were included. Pre-operative, clinical, USG and intra-operative difficulties that were identified beforehand were analysed against the endpoint of difficult cholecystectomy. Weightage was given to each parameter based on the number of times they were found to be significantly associated with each difficulty criteria. As in Randhava *et al.*⁷ study in our study also shows advancing age is the associated with the difficult cholecystectomy and age group of 41-50 years were found for difficult cholecystectomy (48.3%). This study also reveals that male gender, multiple attacks of cholecystitis and previous abdominal surgery is one the the factor for difficult LC as similar to the study of various authors like Kama NA *et al* and others.^{8,9,10} This study also revealed prominently patients with previous ERCP with CBD stent had difficult LC. 10 (83.33%) out of 12 patients had difficulty and even conversion to open cholecystectomy in 2 patients. Another predictive factor for difficult LC was clinically palpable GB due to impacted stone in the Hartman pouch, 11 (78.6%) out of 14 were had difficult cholecystectomy similar to Randhava *et al.* and Agrawal *et al.*^{7,11} Similarly 24(64%) out of 37 of contracted and fibrosed GB found difficult associated with difficulty in dissection at Calot's triangle and separation from GB bed and final extraction. Similar to^{12,13,14,15} This study also observed that distended GB 11(68.7%) out of 14 which was also clinically palpable were found difficult, similar to Singh *et al* study.¹⁶ though 16 patients had distended GB out of which 14 were palpable. Thick wall GB >4 mm 13 (86.7%) out of 15 were found difficult similar to Mohanty SK *et al.*¹⁹ However, in our study we do not found any association in patients with h/o acute cholecystitis, obesity and pericholecystic fluid. All these cases were turned out to be easy LC.

After comparing all the parameters and predictors for difficult cholecystectomy, post ERCP patients were found to be most difficult to perform laparoscopic cholecystectomy. Out of 12 patients with the history of ERCP with CBD stent placement were operated and 10 patients were difficult and 2 patients were converted to open cholecystectomy.

CONCLUSION

Laparoscopic cholecystectomy being established as a gold standard for heavy population presented with symptomatic cholelithiasis, it is utmost important to prepare before hand for difficulty in procedure, avoid complications and legal aspect attached with it. This study demonstrates that preoperative evaluation for predicting the difficulty in LC is feasible.

This study was targeted at identifying and comparing the various possible predictors of difficulty in LC. At the end of this study the following conclusions may be drawn.

- Elderly patients are more likely to have a difficult LC.
- Females undergo this surgery more frequently but males tend to have a higher number of difficult cases.
- Recurrent cholecystitis is a predictor.
- Previous surgery predisposes towards difficulties in cholecystectomy.
- Patients who needed preoperative ERCP had more chances of having a difficult cholecystectomy.
- Abnormal serum hepatic and pancreatic enzyme profiles were associated with difficulty in surgery.
- Pre operative USG can predict difficulties during LG.
- Features like distended or contracted gallbladder, intraperitoneal adhesions, structural anomalies or distortions and the presence of a cirrhotic liver are signs that are associated with subsequent difficulties during the surgery.
- Obese patients and those with acute cholecystitis and pericholecystic fluid tend to have no more difficulties during surgery.

After analysing all the study and difficult level in surgery it is concluded that

1. Most important predictive factor for difficult LC was ERCP with CBD stent of the rate of 83.33%, meaning thereby 10 patients with difficult LC out of total 12 patients of ERCP with CBD stent.
2. Impacted stone at neck of the GB was another important predictive factor found in this study of the rate of 78.8% meaning thereby 17 patients had difficult LC out of total 24 patients with impacted stone at GB neck.
3. This study also showed that conversion to open cholecystectomy in 11 pt out of total 31 was due to 2 patient of ERCP with CBD stent and 10 with impacted stone at neck and 1 one of them having both ERCP stenting as well as impacted stone at neck.
4. Another feature of the study was clinically palpable GB was also predicted as an important factor for difficulty in LC by the rate of 78.0% meaning thereby 11 patient were difficult out of total 14 pt of palpable GB mostly due to impacted stones.
5. During per abdomen examination tenderness in RHC which was observed in 7 patients, out of these 5 (71.4%) patients showed difficulty during LC therefore, it turned out to be a important predictive factor for difficult laparoscopic cholecystectomy.

Reference

1. Hussain A. Difficult laparoscopic cholecystectomy: current evidence and strategies of management. *Surg Laparosc Endosc Percutan Tech.* 2011;21(4):211-217. doi:10.1097/SLE.0b013e318220f1b1
2. Dhanke PS, Ugane SP. Factors predicting difficult laparoscopic cholecystectomy. *Int J Stud Res.* 2014;4:37.
3. Lam CM, Murray FE, Cuschieri A (1996) Increased cholecystectomy rate after the introduction of laparoscopic cholecystectomy in Scotland. *Gut* 38:282–284
4. The southern Surgeons Club, Meyers WC (1991) A prospective analysis of 1518 laparoscopic cholecystectomy. *N Engl J Med* 324:1073–1078
5. Solte M, Radoňak j, A risk score to predict the difficulty of elective laparoscopic cholecystectomy, Department of Surgery, University of Pavol Jozef Safarik, Kosice, Slovak Republic, *Videosurgery Miniinv* 2014; 9
6. Stoikes N., Brunt L.M. (2020) The Difficult Cholecystectomy. In: Asbun H., Shah M., Ceppa E., Auyang E. (eds) *The SAGES Manual of Biliary Surgery*. Springer, Cham. https://doi.org/10.1007/978-3-030-13276-7_11.
7. Randhawa JS, Pujahari AK. Preoperative prediction of difficult lap chole: a scoring method *Indian J Surg.* 2009;71(4):198-201.
8. Kama NA, Kologlu M, Doganay M *et al.* Risk score conversion from laparoscopic to open cholecystectomy. *Am J Surg.* 2001;181:520-5.
9. Hayama S, Ohtaka K, Shoji Y, *et al.* Risk Factors for Difficult Laparoscopic Cholecystectomy in Acute Cholecystitis. *JSLs.* 2016;20(4):e2016.00065. doi:10.4293/JSLs.2016.00065
10. Nidoni R, Udachan TV, Sasnur P, Baloorkar R, Sindgikar V, Narasangi B. Predicting difficult laparoscopic cholecystectomy based on clinicoradiological assessment. *JCDR.* 2015 Dec;9(12):PC09.
11. Agrawal N, Singh S, Khichy S. Preoperative Prediction of Difficult Laparoscopic Cholecystectomy: A Scoring Method. *Niger J Surg.* 2015;21(2):130-33.
12. Sandhu G, Rana M L, Singh K. Preoperative prediction of difficult laparoscopic cholecystectomy: A scoring method. *Ind J of Applied research.* 2016;6(6):481- 87.
13. Hutchinson CH, Traverso LW, Lee FT. Laparoscopic Cholecystectomy. Do preoperative factors predict the need to convert to open? *Surg Endosc.* 1994;8:875-8.
14. Alponat A, Kum CK, Koh BC *et al.* Predictive factors for conversion of laparoscopic cholecystectomy. *World J Surg.* 1997;21(6):629-33
15. Kumar S, Tiwari S, Agrawal N *et al.* Predictive Factors for difficult surgery in laparoscopic cholecystectomy for chronic cholecystitis. *The Internet Journal of surgery.* 2008;16:254-8.
16. Singh K, Ohri A. Difficult laparoscopic cholecystectomy: A large series from north India. *Indian J Surg.* 2006;68:205-8.
17. Mohanty SK, Mohanty R. Preoperative Prediction of difficult Laparoscopic cholecystectomy using clinical and ultrasonographic parameters. *Ann. Int. Med. Den. Res.* 2017;3(4):43-48.