



## Subject Area : Forensic Medicine

# MORPHOMETRIC ANALYSIS OF THE HUMAN STERNUM FOR SEX ESTIMATION: A QUANTITATIVE APPROACH IN FORENSIC MEDICINE

<sup>1</sup>Surya Kiran Panga , <sup>2</sup>Dibya Sharma and <sup>3</sup>Manoj Kumar Wahi

<sup>1</sup>Senior Resident, <sup>2</sup>Senior Resident and <sup>3</sup>Director Professor

<sup>1</sup>Department of Forensic Medicine & Toxicology, AIIMS, New Delhi, India- 110029.

<sup>2</sup>Department of Forensic Medicine, Sikkim Manipal Institute of Medical Sciences, Sikkim, India- 737102.

<sup>3</sup>Department of Forensic Medicine & Toxicology, Lady Hardinge Medical College, New Delhi, India- 110001.

ARTICLE INFO	ABSTRACT
Received 16 <sup>th</sup> March 2024 Received in revised form 23 <sup>rd</sup> March, 2024 Accepted 18 <sup>th</sup> April, 2025 Published online 28 <sup>th</sup> April, 2025	<p><b>Background:</b> Sex estimation is a fundamental step in the identification process during medico-legal investigations, particularly when dealing with fragmented or decomposed remains. The sternum, owing to its resilience and anatomical variability between sexes, serves as a valuable skeletal element for this purpose. <b>Aim:</b> This study aims to evaluate the reliability of various sternal measurements in determining sex in an Indian population. <b>Methods:</b> An observational cross-sectional study was conducted at the Department of Forensic Medicine and Toxicology, VMMC &amp; Safdarjung Hospital, New Delhi, from January 2020 to June 2021. A total of 110 autopsied individuals aged between 10 and 70 years were included. Exclusion criteria comprised cases with fractured, deformed, or diseased sterna. After removal and maceration of the sternum, six osteometric parameters were recorded: length of manubrium, body of sternum, total sternum, and breadths of the manubrium and body, along with the sternal index. Receiver Operating Characteristic (ROC) curve analysis was used to evaluate the discriminative power of each parameter. <b>Results:</b> The study population comprised 84 males (76.36%) and 26 females (23.64%). All sternal parameters, except the sternal index (higher in females), were significantly greater in males (<math>p &lt; 0.0001</math>). ROC analysis revealed that the length of the body of sternum was the most reliable predictor of sex, with an Area Under the Curve (AUC) of 0.906, sensitivity of 85.71%, specificity of 88.46%, and diagnostic accuracy of 86.36%. A cut-off value of <math>&gt;86</math> mm for males and <math>\leq 86</math> mm for females was established. The total sternal length and other parameters also showed acceptable to excellent diagnostic performance. <b>Conclusion:</b> Sternal measurements, particularly the length of the body of the sternum (86.36%), are reliable indicators for sex estimation in forensic settings. This study provides population-specific cut-off values for an Indian cohort, enhancing the utility of sternal morphometry in forensic anthropology and medico-legal practice.</p>
<p><b>Key words:</b></p> <p>Forensic identification, Sternum, Sex estimation, Osteometry, Sternal index, ROC curve, Indian population</p>	
Copyright©	Copyright© The author(s) 2025, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Identification is the process of establishing the individuality of a person, living or deceased. In medico-legal investigations, skeletal examination is essential for identifying unknown individuals, particularly for estimating sex, age, and

stature (1,2). Other features such as race, scars, moles, and professional marks also aid in identification (3). Mass disasters such as plane crashes or fires often result in fragmented or decomposed remains, complicating identification (4). In such cases, forensic experts rely heavily on bones, particularly when soft tissues are destroyed.

The sternum is a flat, elongated bone forming the central anterior thoracic wall, protecting the mediastinal organs, especially the heart (5,6,7). Being subcutaneous and relatively resilient to decomposition, it is often preserved in forensic contexts.

\*Corresponding author: **Dr Surya Kiran Panga**

Department of Forensic Medicine & Toxicology, AIIMS, New Delhi, India- 110029

Sex determination is a critical first step in biological profiling, as it immediately halves the potential pool of identities (8). Reliable sex estimation requires high accuracy (correct identification within a sample) and reliability (performance on independent samples). Legal standards demand at least 95% accuracy (9). Skeletal sex differences become marked after puberty, typically around 15–18 years, with the pelvis, skull, sternum, and long bones providing the most reliable indicators (10,11).

Krogman reported 100% accuracy with complete skeletons, 95% using the pelvis alone, and 80% with long bones. The sternum shows about 80% accuracy in sex estimation (10). Since Wenzel (1788), the sternum has been studied as a single indicator, showing proportionally longer mesosterna in males (12). Dwight found a manubrium-to-body length ratio of 49:100 in males and 52:100 in females (10,13). The sternal index, calculated as (manubrium/body)  $\times$  100, averages 46.2 in males and 54.3 in females (14). Hyrtl's Law states this ratio is  $>1.2$  in females and  $<1.2$  in males (15). Ashley's "149 rule" suggests that a combined manubrium and body length  $\geq 149$  mm indicates male sex, with  $\sim 80\%$  accuracy (16–18).

This study aims to evaluate the sternum for sex estimation, as it often remains intact even in severely decomposed bodies.

## METHODOLOGY

### Study Design and Participants

This study is an prospective observational Cross-Sectional study carried out in the Department of Forensic Medicine and Toxicology, Vardhman Mahavir Medical College & Safdarjung Hospital's, New Delhi, from Jan 2020 to June 2021 after obtaining hospital ethical committee approval (IEC/VMMC/SJH/Thesis/2019-10/157). Dead bodies of Indian population brought for autopsy to the Mortuary wing were the source population.

### Sample Size

In the study of Goksin Nilufer Yonguc et al (19), observed that correlation coefficient of total sternal length and stature was 0.806. Taking this value as reference, the minimum required sample size with 80% power of study and 5% level of significance is 10 study subjects. Also, sensitivity and specificity of total sternal length for predicting gender was 89.2% and 80% respectively. Taking this value as reference, the minimum required sample size with desired precision of 10%, 80% power of study, and 5% level of significance is 108 study subjects. To reduce margin of error, total sample size taken is 110.

### Inclusion & Exclusion Criteria

Cases of both sexes where identity is known and Age greater than 10 years and less than 70 years were included in the study. Whereas, fracture of sternum, Unclaimed dead bodies, Skeletal deformities, diseased or deformed bones that affects sternum were excluded from the study.

### Procedure and technique for the Removal of the Sternum

After obtaining informed consent from the deceased's relative, the body was placed in the supine position with arms by the sides, and an I-shaped incision was made from the chin to the symphysis pubis. The chest muscles were dissected

with the knife directed inward towards the ribs, extending laterally to the mid-axillary line and over the clavicles. The chest cavity was opened by cutting the costal cartilages close to the costochondral junctions, starting from the second cartilage. The sternoclavicular joint was identified by shoulder movement, and its capsule was divided using a vertical, circular motion of the knife. The sternum was detached from the diaphragm and removed completely, with each specimen labelled. Maceration of soft tissues was performed by boiling the sternum in a solution of 7–8 grams of sodium hydroxide per litre of water, using 2 litres of solution in an aluminium vessel for 60–75 minutes. The specimens were monitored to prevent over-boiling and achieve optimal maceration. After boiling, the specimens were thoroughly rinsed, and residual soft tissue was removed using forceps and a soft brush. The cleaned bones were dried on a clean cloth for 2–3 days and then sealed in airtight zip-lock bags for preservation as shown in Figure 1.

### Osteometric Parameters Used for Analysis

Measurements were taken using vernier callipers as shown in Figure 2,

1. Total length of sternum: Straight distance from the deepest point of the suprasternal notch to the point on the lower margin of the corpus Sterna in the mid-sagittal plane. When the sternum was not united as a single piece due to lack of fusion, individual segments were measured independently, and the total length is calculated by adding all.
2. Length of body of sternum: Straight distance from the point on the lower margin of manubrium to the point on the lower margin of the body in the mid-sagittal plane.
3. Length of manubrium: Straight distance from the suprasternal notch to the point on the lower margin of the manubrium in the mid-sagittal plane.
4. Breadth of manubrium: Straight distance between the most laterally placed points on the lateral margins of the manubrium, taken at right angles to the length of the manubrium.
5. Breadth of body of sternum: The straight distance between the most laterally placed points on the lateral margins of the body taken at right angles to the length of the body.
6. Sternal index:

$$\frac{\text{Length of manubrium}}{\text{Length of body}} \times 100$$

### Statistical Analysis


Categorical variables will be presented in number and percentage (%) and continuous variables will be presented as mean  $\pm$  SD and median. Normality of data will be tested by Kolmogorov-Smirnov test. If the normality is rejected, then non parametric test will be used. Pearson correlation coefficient/Spearman Rank correlation coefficient (when the data sets are not normally distributed) will be used to correlate quantitative parameters. Receiver operating characteristic curve will be



used to find out the cut-off point of sternal lengths and breadths for predicting gender. Diagnostic test will be used to calculate sensitivity, specificity, NPV and PPV. A p value of less than 0.05 will be considered as significant. The data will be entered in MS EXCEL spreadsheet and analysis will be done using Statistical Package for Social Sciences (SPSS) version 21.0.

## RESULTS AND OBSERVATIONS

In the present study, 26.36% of patients belonged to age group of 31-40 years followed by 21-30 years (23.64%), 41-50 years (21.82%), 51-60 years (12.73%), 11-20 years (8.18%) and 61-70 (7.27%). Mean value of age(years) of study subjects was  $38.22 \pm 13.7$  with median (25th-75th percentile) of 35.5(29-47.75) as shown in figure 3.




**Figure 1.** Labelled and Packed in Air Tight Zip Locker

In present study, 76.36% of patients were males and 23.64% of patients were females.

Mean value of length of manubrium (mm), length of body of sternum (mm), length of total sternum (mm), breadth of manubrium (mm), breadth of body of sternum (mm) and sternal index of study subjects was  $163.29 \pm 8.16$ ,  $51.43 \pm 5.19$ ,  $90.99 \pm 11.77$ ,  $166.46 \pm 19.24$ ,  $57.37 \pm 7.25$ ,  $37.24 \pm 6.47$  and  $57.49 \pm 8.07$  with median (25th-75th percentile) of 163(158-169), 51(48-54.95), 91.85(84.5-99.375), 167.5(154-180.375), 57(52-61), 37(32-42) and 57.06(52.65-62.092) respectively as shown in table 1.

ROC curves above the diagonal line are considered to have reasonable discriminating ability to predict sexual dimorphism. All the parameters had significant discriminatory power to predict male. Interpretation of the area under the ROC curve showed that the performance of length of body of sternum (mm) (AUC 0.906; 95% CI: 0.836 to 0.954) was outstanding. Discriminatory power of length of total sternum (mm) (AUC 0.894; 95% CI: 0.821 to 0.945) was excellent and discriminatory power of length of manubrium (mm) (AUC 0.75; 95% CI: 0.658 to 0.827), breadth of manubrium (mm) (AUC 0.782; 95% CI: 0.693 to 0.855), breadth of body of sternum (mm) (AUC 0.769; 95% CI: 0.679 to 0.844) and sternal index (AUC 0.739; 95% CI: 0.646 to 0.818) was acceptable. Among all the parameters, Length of body of sternum (mm) was the best predictor of male at cut off point of >86 with 90.60% chances of correctly predicting gender sex as male.



**Figure 2.** Osteometric Measurements By Vernier Callipers.

Length of body of sternum (mm) had sensitivity of 85.71% followed by breadth of body of sternum (mm) (84.52%), length of total sternum (mm) (82.14%), breadth of manubrium (mm) (79.76%). In prediction of male, Length of manubrium (mm) had lowest sensitivity of 52.38%. On the other hand, length of body of sternum (mm), length of manubrium (mm) had specificity of 88.46% each followed by length of total sternum (mm) (84.62%), breadth of body of sternum (mm)

**Table 1.** Descriptive statistics of measurements of study subjects.

Measurements	Mean $\pm$ SD	Median (25th-75th percentile)	Range
Length of manubrium (mm)	$51.43 \pm 5.19$	51(48-54.95)	37.5-67.5
Length of body of sternum (mm)	$90.99 \pm 11.77$	91.85(84.5-99.375)	66-119
Length of total sternum (mm)	$166.46 \pm 19.24$	167.5(154-180.375)	115.5-216
Breadth of manubrium (mm)	$57.37 \pm 7.25$	57(52-61)	40-83.5
Breadth of body of sternum (mm)	$37.24 \pm 6.47$	37(32-42)	24-56
Sternal index	$57.49 \pm 8.07$	57.06(52.65-62.092)	39.61-85.07

**Table 2.** Receiver operating characteristic curve of measurements to predict sex as male.

Male	Length of manubrium (mm)	Length of body of sternum (mm)	Length of total sternum (mm)	Breadth of manubrium (mm)	Breadth of body of sternum (mm)	Sternal index
Area under the ROC curve (AUC)	0.75	0.906	0.894	0.782	0.769	0.739
Standard Error	0.0476	0.0342	0.0373	0.0488	0.059	0.0543
95% Confidence interval	0.658 to 0.827	0.836 to 0.954	0.821 to 0.945	0.693 to 0.855	0.679 to 0.844	0.646 to 0.818
P value	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
Cut off	>51.5	>86	>159	>52	>32	≤60.5
Sensitivity(95% CI)	52.38%(41.2 - 63.4%)	85.71%(76.4 - 92.4%)	82.14%(72.3 - 89.6%)	79.76%(69.6 - 87.7%)	84.52%(75.0 - 91.5%)	79.76%(69.6 - 87.7%)
Specificity(95% CI)	88.46%(69.8 - 97.6%)	88.46%(69.8 - 97.6%)	84.62%(65.1 - 95.6%)	61.54%(40.6 - 79.8%)	65.38%(44.3 - 82.8%)	65.38%(44.3 - 82.8%)
PPV(95% CI)	93.6%(82.5 - 98.7%)	96%(88.8 - 99.2%)	94.5%(86.6 - 98.5%)	87%(77.4 - 93.6%)	88.7%(79.7 - 94.7%)	88.2%(78.7 - 94.4%)
NPV(95% CI)	36.5%(24.7 - 49.6%)	65.7%(47.8 - 80.9%)	59.5%(42.1 - 75.2%)	48.5%(30.8 - 66.5%)	56.7%(37.4 - 74.5%)	50%(32.4 - 67.6%)
Diagnostic accuracy	60.91%	86.36%	82.73%	75.45%	80.00%	76.36%

**Table 3.** Receiver operating characteristic curve of measurements to predict sex as female.

Female	Length of manubrium (mm)	Length of body of sternum (mm)	Length of total sternum (mm)	Breadth of manubrium (mm)	Breadth of body of sternum (mm)	Sternal index
Area under the ROC curve (AUC)	0.75	0.906	0.894	0.782	0.769	0.739
Standard Error	0.0476	0.0342	0.0373	0.0488	0.059	0.0543
95% Confidence interval	0.658 to 0.827	0.836 to 0.954	0.821 to 0.945	0.693 to 0.855	0.679 to 0.844	0.646 to 0.818
P value	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
Cut off	≤51.5	≤86	≤159	≤52	≤32	>60.5
Sensitivity(95% CI)	88.46%(69.8 - 97.6%)	88.46%(69.8 - 97.6%)	84.62%(65.1 - 95.6%)	61.54%(40.6 - 79.8%)	65.38%(44.3 - 82.8%)	65.38%(44.3 - 82.8%)
Specificity(95% CI)	52.38%(41.2 - 63.4%)	85.71%(76.4 - 92.4%)	82.14%(72.3 - 89.6%)	79.76%(69.6 - 87.7%)	84.52%(75.0 - 91.5%)	79.76%(69.6 - 87.7%)
PPV(95% CI)	36.5%(24.7 - 49.6%)	65.7%(47.8 - 80.9%)	59.5%(42.1 - 75.2%)	48.5%(30.8 - 66.5%)	56.7%(37.4 - 74.5%)	50%(32.4 - 67.6%)
NPV(95% CI)	93.6%(82.5 - 98.7%)	96%(88.8 - 99.2%)	94.5%(86.6 - 98.5%)	87%(77.4 - 93.6%)	88.7%(79.7 - 94.7%)	88.2%(78.7 - 94.4%)
Diagnostic accuracy	60.91%	86.36%	82.73%	75.45%	80.00%	76.36%

**Table 4.**Comparative statistics of manubrial and body of sternal lengths, mm (mean $\pm$ 3SD) of sternum studied by different workers (21-22, 25-34).

Author	Region	Sex	N	Length of Manubrium	t-test	Length of body of sternum	t-test
Dwight et al.	-	M	142	53.70	-	110.40	-
		F	86	49.40	-	91.90	-
Paterson et al.	-	M	310	52.00	-	103.70	-
		F	126	47.30	-	91.00	-
Ashley et al.	Africa	M	85	45.90	-	96.50	-
		F	13	44.20	-	82.90	-
Ashley et al.	Europe	M	378	52.20	-	104.70	-
		F	168	47.90	-	90.80	-
Selthofer et al.	Croatia	M	55	55.2 $\pm$ 3.6	-5.25**	109.7 $\pm$ 14.4	-7.59**
		F	35	52.4 $\pm$ 4.5	-5.55**	94.2 $\pm$ 14.0	-5.95*
Jit et al.	Chandigarh (India)	M	312	51.73 $\pm$ 5.22	0.83	95.35 $\pm$ 9.88	-1.52
		F	88	48.42 $\pm$ 5.28	-1.60	78.60 $\pm$ 9.31	-0.04
Dahiphale et al.	Maharashtra (India)	M	96	48.46 $\pm$ 5.59	5.50**	94.43 $\pm$ 9.43	-0.31
		F	47	43.78 $\pm$ 5.24	3.60**	70.19 $\pm$ 8.54	5.03**
Gautam et al.	Gujarat (India)	M	56	53.00	-	95 . 00	-
		F	44	48.00	-	76.00	-
Torwalt et al.	Canada	M	89	54.43 $\pm$ 6.10	-3.20**	109.41 $\pm$ 12.48	- 10.42**
		F	41	48.79 $\pm$ 4.92	-1.71	93.55 $\pm$ 8.99	8.41**
Queiroz et al.	Brazil	M	50	-	-	88 .87 $\pm$ 10.04	3.32**
		F	50	-	-	82.44 $\pm$ 10 .05	-2.17*
Vella et al.	Italy	M	20	52.60 $\pm$ 5.50	-0.38	97.7 $\pm$ 7.70	-1.94
		F	20	43.20 $\pm$ 4.90	3.18**	85.70 $\pm$ 10.30	-2.76*
Fernandez et al.	Spain	M	36	51.56 $\pm$ 5.63	0.54	101.13 $\pm$ 7.69	-4.59**
		F	40	45.00 $\pm$ 5.51	2.10*	82.71 $\pm$ 8.40	-2.32*
Hunnargi et al.	Maharashtra (India)	M	75	51.99 $\pm$ 4.96	0.17	89.17 $\pm$ 10.63	1.32
		F	40	44.88 $\pm$ 10.63	3.53**	72.38 $\pm$ 9.85	3.63**
Atal et al.	Delhi (India)	M	50	45.75 $\pm$ 2.99	11.76**	100.28 $\pm$ 6.25	-5.68**
		F	50	41.20 $\pm$ 3.31	8.32**	78.35 $\pm$ 6.26	0.14
Present study (2021)	Delhi (India)	M	84	52.42 $\pm$ 5.21	-	94.92 $\pm$ 9.65	-
		F	26	48.23 $\pm$ 3.65	-	78.27 $\pm$ 8.62	-

\*\*Significant at 0.01 level. \*Significant at 0.05 't'-Values for comparison with present study.

(65.38%). In prediction of male, Breadth of manubrium (mm) had lowest specificity of 61.54%. Highest positive predictive value was found in length of body of sternum (mm) (96.00%) and highest negative predictive value was found in length of body of sternum (mm) (65.70%). There is always a trade-off between sensitivity and specificity (any increase in sensitivity will be accompanied by a decrease in specificity) so we choose that variable as best in which combination of sensitivity and specificity gives the maximum predictive value i.e., maximum diagnostic accuracy. So overall length of body of sternum (mm) was best predictor of gender as shown in table 2.

ROC curves above the diagonal line are considered to have reasonable discriminating ability to predict female. All the parameters had significant discriminatory power to predict female. Interpretation of the area under the ROC curve showed that the performance of length of body of sternum

(mm) (AUC 0.906; 95% CI: 0.836 to 0.954) was outstanding. Discriminatory power of length of total sternum (mm) (AUC 0.894; 95% CI: 0.821 to 0.945) was excellent and discriminatory power of length of manubrium (mm) (AUC 0.75; 95% CI: 0.658 to 0.827), breadth of manubrium (mm) (AUC 0.782; 95% CI: 0.693 to 0.855), breadth of body of sternum (mm) (AUC 0.769; 95% CI: 0.679 to 0.844) and sternal index (AUC 0.739; 95% CI: 0.646 to 0.818) was acceptable. Among all the parameters, Length of body of sternum (mm) was the best predictor of female at cut off point of  $\leq 86$  with 90.60% chances of correctly predicting female.

Length of manubrium (mm) and length of body of sternum (mm) had sensitivity of 88.46% each followed by length of total sternum (mm) (84.62%), breadth of body of sternum (mm) (65.38%). In prediction of female, Breadth of manubrium (mm) had lowest sensitivity of 61.54%. On the other hand, length



of body of sternum (mm) had specificity of 85.71% followed by breadth of body of sternum (mm) (84.52%), length of total sternum (mm) (82.14%), sternal index (79.76%). In prediction of female, Length of manubrium (mm) had lowest specificity of 52.38%. Highest positive predictive value was found in length of body of sternum (mm) (65.70%) and highest negative predictive value was found in length of body of sternum (mm) (96.00%). So we choose that variable as best in which combination of sensitivity and specificity gives the maximum predictive value i.e. maximum diagnostic accuracy so overall length of body of sternum (mm) was best predictor of female as shown in table 3.

## DISCUSSION & CONCLUSION

It was obvious from the foregoing observations that all parameters measured have a higher value in males except sternal index which is greater in females. This is in total agreement with previously studies (10, 20-24). Table 4 shows the comparative statistics of manubrial and body of sternal lengths, mm (mean $\pm$ 3SD) of sternum studied by different workers in various parts of world including India.

All the parameters of the sternum i.e., the length, breadth of the body, breadth of the manubrium, sternal index and total length showed significant statistical difference between males and females. The sternal index is found characteristically high in all the studies in females (10, 14, 35, 36). This is reflected in the higher mean values obtained for females ( $62.11 \pm 6.67$ ) in this study, when compared to that of males ( $56.06 \pm 7.95$ ). Sternal index is calculated from the formula: Length of manubrium X 100/ Length of body. The reason for the higher value of sternal index in females may be attributed to the following: The length of manubrium exceeds half the length of body of the sternum in females; while in males, the length of body of sternum is at least twice the length of manubrium. This view which had been put forward by many authors and workers has also been noticed in the present study (35, 36). This observation could also explain the reason for the statistically significant difference in the measurements of length of manubrium in males and females. One author is of the view that the manubrium is somewhat smaller in males while bigger in females (37). Jit et al., have opined that manubrium corpus index is not useful in sexing a given sternum (23). Dwight et al., stated that variations in the length of manubrium in the two sexes are minimal but an individual sternum is usually found to lie in the overlapping zone (26). Only a negligible percentage of sternums could be sexed accurately in the present study, as well as by Jit et al., using demarking points for length of manubrium (22). Using limiting point for this variable while the present study could estimate sex with an accuracy of 60.91%, whereas in study by Hunnargi et al., this variable achieved an accuracy of around 77% (29).

The mean length of body of the sternum was found to be significantly higher in males (94.92mm) than females (78.27 mm). This observation is in close agreement with those of Wenzel et al., Krogman et al., Paterson et al., and Jit et al., have all suggested that length of body of the sternum is greater in males (10, 20-22). The accuracy for sex estimation based on limiting point for length body of the sternum in the present study (86.36%) closely match that found by Hunnargi et al (29).

The mean total length of sternum is also significantly higher

in males (172.7mm) than in females (146.31mm). This finding again is in concordance with the observation of Jit et al (22). The accuracy of sex estimation derived from combined length of manubrium and sternum using either Ashley's 'rule 149' (by Jit et al., and Hunnargi et al.) or using limiting point (by Hunnargi et al.; and present study) happens to be almost similar ranging between 82.1 and 94.5% and was in concordance with the present study (accuracy: 82.73%) (22, 29).

ROC curves above the diagonal line are considered to have reasonable discriminating ability to estimate sex. All the parameters had significant discriminatory power to predict male. Length of body of sternum (mm) had sensitivity of 85.71% and length of manubrium (mm) had lowest sensitivity of 52.38%. On the other hand, both the length of body of sternum (mm) and the length of manubrium (mm) had specificity of 88.46% and breadth of manubrium (mm) had lowest specificity of 61.54%. Present study had mentioned a definite cut off value above or below which the sternum can be sexed. Therefore, a subject is likely **male** if:

- Sternal body > 86 mm
- Manubrium length > 51.5 mm
- Sternal body breadth > 32 mm
- Manubrium breadth > 52 mm
- Sternal index  $\leq$  60.5
- Total sternum length > 159 mm

Likely **female** if these values are equal to or below the above thresholds, where as , it vice versa for sternal index.

There is always a trade-off between sensitivity and specificity (any increase in sensitivity will be accompanied by a decrease in specificity) so we choose that variable as best in which combination of sensitivity and specificity gives the maximum predictive value i.e., maximum diagnostic accuracy. So, overall the length of body of sternum (86.36%) was best predictor of Sex.

## References

1. Reddy KSN, Murty OP. The essentials of Forensic Medicine & Toxicology. 34th ed. New Delhi: Jaypee; 2017. p62-76.
2. Aggrawal A. Textbook of Forensic Medicine and Toxicology. 2<sup>nd</sup> ed. New Delhi: APC; 2021. p73.
3. Manoharan C, Dhanalakshmi V, Thangam D, Joe AE. Estimation of age from human sternum-an autopsy study. *Indian J Forensic Community Med.* 2016;3(2):128-32.
4. Manoharan C, Jeyasingh T, Dhanalakshmi V, Thangam D. Is Human Sternum a Tool for Determination of Sex?. *Indian J Forensic Community Med.* 2016;3(1):60-63.
5. Snell RS. Clinical anatomy for Medical students. 3rd ed. Boston: Little Brown and Company; 1991. p35-36.
6. Moore KL, Dalley AF, Agur AMR. MOORE Clinically oriented anatomy. 7th ed. Philadelphia: LWW; 2015. p76-78.
7. Soames RW. Gray's Anatomy. 38th ed. New York: ELBS with Churchill Livingstone; 1995. p537-539.
8. Knight B. Forensic pathology. 2<sup>nd</sup> ed. London: Arnold publication; 1996. p. 113.
9. Schmitt A, Cunha E, Pinheiro J. Forensic anthropology and medicine. 1st ed. New Jersey: Humana; 2006. p225-

- 226.
10. Krogman WM. Human Skeleton in Forensic Medicine. Illinois: Charles C. Thomas; 1962. p215-217.
11. Camps FE, Gradwohl RBH. Gradwohl's Legal Medicine. 2<sup>nd</sup> ed. Bristol: J Wright; 1968. p127.
12. Krogman WM. Human Skeleton in Forensic Medicine. Illinois: Charles C. Thomas; 1962. p215-217.
13. Stewart TD. Essentials of Forensic Anthropology. 1<sup>st</sup> ed. Illinois: Charles C Thomas; 1979. p95.
14. Smith S, Fiddes FS. Forensic Medicine - A textbook for students and practitioners. 10<sup>th</sup> ed. London: J and A Churchill Ltd; 1955. p69-77.
15. Hyrtl J. Handbuch der Topographischen Anatomic percentage. 1<sup>st</sup> ed. Vienna: Wilhelm Braumuller; 1893. p348.
16. Ashley GT. Typing of the human sternum. The influence of sex and age on its measurements. *J of forensic med.* 1956;3:27-43.
17. Ashley. Frazer's Anatomy of Human skeleton. 5<sup>th</sup> ed. London: Churchill; 1956. p56, 58,59.
18. Rentoul E, Smith H. Medical Jurisprudence and Toxicology. 13<sup>th</sup> ed. Edinburgh: E and S Livingstone; 1973. p80.
19. Yonguc GN et al. Estimation of stature and sex from sternal lengths: an autopsy study. *Anat Sci Int.* 2015;90:89-96.
20. Wenzel J. A Comparison of human and anthropoid mesosterna. *Am J Phys Anthropol.* 1788;3:449-461.
21. Paterson A.M. The human sternum. Liverpool: Williams and Norgate; 1904. p36,37,77.
22. Jit I, Jhingan V, Kulkarni M. Sexing the human sternum. *Am J Phys Anthropol.* 1980;53(2):217-24.
23. Jit I, Kaur H. Time of fusion of the human sternbrae with one another in North West India. *Am J of Phys Anthropol.* 1989;80(2):195-202.
24. Buikstra JE, Ubelaker DH. Standards for data collection from human skeletal remains. Fayetteville: Arkansas Archeological Survey Research Series No. 44; 1994.
25. Dahiphale VP, Baheete BH. Sexing the human sternum in Marathwada Region. *J Anat Soc India.* 2002;51(2):162-167.
26. Dwight T. The sternum as an index of sex, age and height. *J of Anat.* 1890;24(4):527- 35.
27. Gautam RS, Shah GV, Jadav HR, Gohil BJ. The human sternum as an index of age & sex. *J Anat Soc India.* 2003;52:20-23.
28. Atal DK, Murari A, Naik SK. Gender differentiation from sternal width. *J Indian Acad Forensic Med.* 2008;30(4):198-201.
29. Hunnargi S, Menezes R, Kanchan T, Lobo S, V S B, Uysal S et al. Sexual dimorphism of the human sternum in a Maharashtrian population of India: A morphometric analysis. *Leg Med.* 2008;6:6-10.
30. Selthofer R, Nicolice V, Mrcela T, Radic R, Leksan I, Rudez I, et al. Morphometric analysis of the sternum. *Coll Antropol.* 2006;30(1):43-47.
31. Torwalt CRMM, Hoppa RD. A test of sex determination from measurements of chest radiographs. *J Forensic Sci.* 2005;50(4):785-790.
32. Queiroz A, Wada RB, Preza A, Spadacio C, Cruz BDL. Sex determination by weight verification of sternum bone Brasil. *Med. leg. Costa Rica.* 2005;22(1):73-78.
33. Vella GD, Campobasso CP, Dragone M, Introna Jr F. Sex determination by sternal measurements on a contemporary Southern Italian population. In: Bergamini PR (ed.). Proceedings of IVth International Meeting on Forensic Medicine. Gorgia: Tipographia, Usl/2 Gorziana;1994. p127-33.\
34. Fernandez ED, Saez AS, Moro JIM. Anthropological determination of sex by studying the sternum. *Rev Esc deMed Leg.* 2007;6:27-42.
35. Polson CJ, Gee DJ, Knight B. Essentials of Forensic Medicine, 3<sup>rd</sup> ed. New York: Pergamon Press; 1973. p48,51.
36. Simpson K. Modern Trends in Forensic Medicine. 1<sup>st</sup> ed. London: Butterworth and Co; 1953. p139.
37. Mukherjee JB. Forensic Medicine and Toxicology. 1<sup>st</sup>ed. Kolkata: Academic Publishers; 1981. p83.

#### How to cite this article:

Surya Kiran Panga., Diba Sharma and Manoj Kumar Wahi. (2025). Morphometric Analysis of the Human Sternum for Sex Estimation: A Quantitative Approach in Forensic Medicine, International Journal of Current Advanced Research, 14(04), pp.196-202.

\*\*\*\*\*